

View this article online at: [patient.info/heart-health/varicose-veins-leaflet](https://patient.info/heart-health/varicose-veins-leaflet)

## Varicose Veins

Studies suggest that varicose veins occur in 10-20 out of 100 men and 25-33 out of 100 women. Most people with varicose veins do not have an underlying disease. They often occur for no apparent reason, although they are associated with lifestyle factors in some cases.

Varicose veins do not cause symptoms or complications in most cases, although some people find them unsightly. If treatment is advised, or wanted for cosmetic reasons, a procedure to seal them off is used. There are several procedures available: heat, lasers or chemicals injected into the veins. These methods have largely replaced the old-fashioned surgical methods such as stripping the veins out.

### What are varicose veins?


Varicose veins are enlarged (dilated) sections of veins which are located just under the surface of the skin - usually on the leg. They are often easy to see, as they look thick and knobbly. They may be less obvious if you are overweight, as they are hidden by fatty tissue under the skin.




By self, Public domain, via Wikimedia Commons

Other, smaller types of veins which can be noticeable are:

- Reticular veins - a closely grouped network of small veins.
- Thread veins, or spider veins - these look like a kind of star burst pattern on an area of the leg. They are not true varicose veins.






#### Healthy venous valves

Venous blood flows upward against gravity and any backflow is prevented by valves that shut against the flow

## What causes varicose veins?



#### Varicose veins

The valves become damaged and do not function properly. Backflow of blood is not prevented and 'pooling' of blood stretches and balloons the vein walls.

It is thought that the wall of the vein becomes weak in some sections. These sections then widen and become more prominent.

If this occurs near a valve then the valve may become leaky and blood may flow backwards. Once this happens at one valve there is extra pressure on the vein. This can cause more widening and more leaky valves. Blood then pools (collects) in the enlarged vein and makes it stand out.

About 10-20 out of 100 men and 25-33 out of 100 women develop varicose veins at some time in their lives. More women than men develop varicose veins. Most people with varicose veins do not have an underlying disease and they occur for no apparent reason. However, the chance of them developing is increased with:

- **Pregnancy.** This is partly due to the baby causing extra pressure on the veins and partly because hormones you make during pregnancy tend to relax vein walls. The more babies you have, the more the risk of permanent varicose veins developing. Varicose veins tend to appear, or become worse, during pregnancy but often improve after childbirth when the pressure on the veins eases.
- **Age.** They are more common with increasing age.
- **Being overweight,** which appears to increase the chance for women but not for men.
- **Standing a lot.** Jobs which involve lots of standing are often said to cause varicose veins. However, there is little scientific evidence to support this theory.

Sometimes an underlying disease may cause varicose veins - for example:

- A previous **blood clot (thrombosis)** or **injury in a deep leg vein.**
- Rarely, a swelling or tumour in the lower part of the tummy (the pelvis) which blocks flow in the veins at the top of the leg.
- Very rarely, varicose veins are due to abnormal blood vessels - if some of the veins or arteries have not formed in the normal way.

## What are the symptoms of varicose veins?

Most people with varicose veins have no symptoms. Some people are concerned about the appearance of the veins. Larger varicose veins can ache, feel heavy or itch.

## Are there any complications of varicose veins?

Most people with varicose veins do not develop complications. Complications develop in a small number of cases. Complications are due to the higher pressure in the varicose veins causing changes to the small blood vessels in nearby skin.

Possible complications include:

- [Inflammation of the vein \(thrombophlebitis\)](#).
- [Swelling of the foot or lower leg](#).
- Skin changes over the prominent veins. The possible skin changes are discolouration, [varicose eczema](#), skin ulcers, or lipodermatosclerosis - hardening of the fat layer under the skin, causing areas of thickened, red skin. [See the separate leaflet called Venous Leg Ulcers for more details](#).
- Rarely, varicose veins may bleed.

**First aid for bleeding varicose veins:** bleeding happens only rarely. If a varicose vein does bleed then you need to stop the bleeding quickly by doing the following:

- Raise the leg - lie down flat and raise the leg high. This will ensure that it is well above the rest of the body (for example, rest it on a chair, use lots of pillows or someone can hold the leg high).
- Pressure - put a clean cloth or dressing on to the bleeding area and put firm pressure on it, for at least 10 minutes.
- Call an ambulance if the bleeding is heavy, or does not quickly stop. See a doctor urgently, as you may need treatment to prevent the bleeding from happening again.

## Treatment

Most people with varicose veins do not need any treatment. You may want to have treatment for one of the following reasons:

- **If complications develop** - these occur in a small number of cases. If leg swelling or skin changes develop over prominent veins, treatment is usually advised to prevent a skin ulcer from developing. If a skin ulcer does occur then treatment of any varicose veins may help to cure the ulcer. If you have a varicose vein which has bled, you should be referred urgently for treatment.
- **For symptoms** of itch or discomfort.
- **Cosmetic reasons.** You may feel that the veins look unsightly. Treatment which is purely for cosmetic reasons is not usually available on the NHS.
- **Combination symptoms:** you will need to have the leg circulation examined before certain treatments (such as compression stockings) can be used if you have a combination of BOTH:
  - Varicose veins which are problematic; AND
  - Arterial disease (poor circulation, or peripheral arterial disease) of the legs.

## Tests

If varicose veins are problematic, you will usually be referred for assessment by a doctor who is a specialist. You may have a type of [ultrasound scan called a Doppler or a duplex scan](#). This helps to show how the blood is flowing in the veins. It can show whether any of the valves are damaged - which is useful to know when planning treatment. Occasionally, other tests are needed if the veins are complex.

If you have arterial disease (poor circulation, or peripheral arterial disease) in your legs, or if arterial disease is suspected, the arterial circulation needs to be measured. This needs to be done before you have treatment which puts pressure on the leg, such as compression stockings (below). The arterial circulation is normally measured by using an ultrasound machine called a Doppler ultrasound, which is used to give a measurement called the ankle brachial pressure index. This test can be done in specialised clinics and also by some nurses and GPs.

## Treatment options

There are several different options. Traditional operations such as vein stripping have largely been replaced by procedures which involve heat, lasers or the injection of chemicals into the vein.

**Self-help methods.** Avoid prolonged standing or sitting still. Try to put your feet up frequently (sit or lie down and raise the feet above the level of your hips). You can, for example, use extra pillows under your feet on a bed or footrest). This helps to reduce blood pooling in the veins. Use a moisturising cream or ointment to protect the skin in the affected area if it is dry, flaky or itchy.

Varicose vein treatment techniques have been developed which have a lower rate of complications, such as bruising, compared with traditional surgery. The National Institute for Health and Care Excellence (NICE) recommends that they be used in preference to traditional surgery for people who are having their veins treated for the first time. They include:

- **Radiofrequency ablation and endovenous laser ablation.** These methods involve passing a probe into one of the longer varicose veins, using ultrasound to guide the position. The laser or radiofrequency energy makes the vein heat up, which seals it.
- **Foam sclerotherapy.** This technique is used if heat or lasers do not work. It uses a chemical mixed with air to make foam. The foam is injected into the veins, pushing the blood away and making the veins go into spasm. After treatment, compression stockings are needed, and the veins will be hard and swollen for a while before they shrink down. More than one treatment may be needed

**Surgery.** Traditional surgery is recommended if treatment with heat, lasers or foam does not work. Different techniques can be used to remove the veins, depending on their site and severity.

**Sclerotherapy.** The vein is injected with a chemical that can close and seal (sclerose) it. It was once used as a main treatment but it became clear that it commonly causes skin staining and ulcers. It is now mainly reserved for small veins below the knee which have not been treated properly or have come back after surgery. The vein needs to be compressed afterwards, which involves wearing bandaging or compression stockings for a few days or weeks.

**Support tights and compression stockings.** These counter the extra pressure in the veins. They may help to ease symptoms such as ache, although there is little proof as to how well they work. They may be difficult to put on, particularly by people who have arthritis in their hands. Current guidelines do not recommend that they be used routinely unless treatments to seal the veins are not suitable or do not work. If you do need compression stockings, below-knee class 1 (light) or class 2 (medium) are usually the most suitable. Ideally, they should be put on first thing in the morning, before you get out of bed, then taken off when going to bed at night. Compression stockings are available on prescription or you can buy them.

**Note:** if you have arterial disease in the legs, you will need a medical assessment of your circulation to decide if compression stockings are suitable (see above).

## Further reading & references

- [Varicose veins: diagnosis and management](#); NICE Clinical Guideline (July 2013)
- [Ultrasound-guided foam sclerotherapy for varicose veins](#); NICE Interventional Procedure Guidance, February 2013
- Kuhlmann A, Prenzler A, Hacker J, et al; Impact of radiofrequency ablation for patients with varicose veins on the budget of the German statutory health insurance system. *Health Econ Rev.* 2013 Apr 3;3(1):9. doi: 10.1186/2191-1991-3-9.
- Kanwar A, Hansrani M, Lees T, et al; Trends in varicose vein therapy in England: radical changes in the last decade. *Ann R Coll Surg Engl.* 2010 May;92(4):341-6.
- Dindyal S, Woodburn KR; Changing practice from conventional surgery to endovenous treatments produces excellent results for both. *Ann R Coll Surg Engl.* 2010 Jan;92(1):87.
- [Varicose veins](#); NICE CKS, February 2020 (UK access only)

**Disclaimer:** This article is for information only and should not be used for the diagnosis or treatment of medical conditions. Egton Medical Information Systems Limited has used all reasonable care in compiling the information but makes no warranty as to its accuracy. Consult a doctor or other healthcare professional for diagnosis and treatment of medical conditions. For details see our [conditions](#).

Author: Dr Laurence Knott	Peer Reviewer: Dr Hayley Willacy	
Document ID: 4357 (v44)	Last Checked: 19/02/2021	Next Review: 18/02/2026

View this article online at: [patient.info/heart-health/varicose-veins-leaflet](http://patient.info/heart-health/varicose-veins-leaflet)

Discuss Varicose Veins and find more trusted resources at [Patient](#).



Book appointments,  
order repeat prescriptions and  
view your medical record online

To find out more visit  
[www.patientaccess.com](http://www.patientaccess.com)  
or download the app

